Jornal de Pediatria xxxx;xxx(xxx): 101463



Jornal de Pediatria

www.jped.com.br



REVIEW ARTICLE

Transition to adolescence in neurodevelopmental disorders: challenges and clinical perspectives

Q1 Eduardo Jorge Custódio da Silva 🌔 ^{a,b,c,d,e,*}, Evelyn Eisenstein 🕞 ^{d,e,f}

Received 12 September 2025; accepted 13 September 2025 Available online xxx

KEYWORDS

Autism spectrum disorder (ASD); ADHD; Neurodevelopment; Adolescence; Adolescent health

Abstract

Objective: This review article describes the main clinical, emotional, educational, and social aspects involved in the transition to adolescence for individuals with these conditions. It also addresses implications for healthcare, the continuity of educational support, and the importance of individualized transition strategies, highlighting the need for a multidisciplinary approach centered on the individual and on continuity of care.

Data sources: Review of the latest literature on neurodevelopmental disorders in adolescents. Data summary: Adolescence represents a vital period of biopsychosocial transformation, particularly challenging for individuals with neurodevelopmental disorders (NDDs), such as autism spectrum disorder (ASD), attention-deficit/hyperactivity disorder (ADHD), and intellectual disability.

Conclusions: Integrated, person-centered, continuous, and collaborative health care is essential to promote the well-being and inclusion of these adolescents.

© 2025 The Authors. Published by Elsevier España, S.L.U. on behalf of Sociedade Brasileira de Pediatria. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

1 Introduction

- Neurodevelopmental disorders (NDDs) are childhood-onset
 conditions characterized by deficits in cognitive, behavioral,
 - * Corresponding author. E-mail: eduardo.jcs@uol.com.br (E.J. Silva).

and social development, including autism spectrum disorder 4 (ASD), attention deficit disorder (ADHD), language and 5 learning disorders, and intellectual disability [1].

The number of cases of children and adolescents with 7 NDDs, also known as neurodivergent disorders, diagnosed in 8 Brazil and worldwide has increased significantly over the 9 past few decades. The causes of this increase are still not 10

https://doi.org/10.1016/j.jped.2025.101463

0021-7557/© 2025 The Authors. Published by Elsevier España, S.L.U. on behalf of Sociedade Brasileira de Pediatria. This is an open access article under the CC BY license (http://creativecommons.org/licenses/by/4.0/).

Please cite this article in press as: E.J. Silva and E. Eisenstein, Transition to adolescence in neurodevelopmental disorders: challenges and clinical perspectives, Jornal de Pediatria (2025), https://doi.org/10.1016/j.jped.2025.101463

^a Universidade Federal do Rio de Janeiro (UFRJ), Neurologia, Rio de Janeiro, RJ, Brazil

^b Instituto Nacional de Saúde da Mulher, da Criança e do Adolescente Fernandes Figueira (IFF/FIOCRUZ), Saúde da Criança, Rio de Janeiro, RJ, Brazil

^c Faculdade de Ciências Médicas da Universidade do Estado do Rio de Janeiro (FCM-UERJ), Departamento de Pediatria, Rio de Janeiro, RJ, Brazil

^d Sociedade Brasileira de Pediatria (SBP), GT sobre Saúde na Era Digital, Rio de Janeiro, RJ, Brazil

^e Academia de Medicina do Rio de Janeiro (AMRJ), Rio de Janeiro, RJ, Brazil

^f Clínica de Adolescentes e do Centro de Estudos Integrados da Infância, Adolescência e Saúde, Rio de Janeiro, RJ, Brazil

E.J. Silva and E. Eisenstein

fully understood; however, it is known that much of it is due to the increased knowledge of these conditions among professionals in various health fields, allowing for the correct diagnosis of previously unrecognized cases and the identification of milder cases.

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

These disorders are currently the most well-known and highly visible conditions that affect the brain and mental and behavioral development during childhood and adolescence. Today, few people have never heard of someone with these disorders or don't have a history of someone with them. In this context, understanding neurodevelopmental disorders is crucial for care for neurodivergent individuals to be based on solid scientific evidence. The vulnerability of these populations, coupled with families' anguish in seeking quick solutions, can lead to adherence to therapies without pediatric technical support. Using a treatment without scientific evidence can lead to significant losses - not just financial ones — because, when deciding to pursue alternative approaches, many families end up abandoning interventions with proven efficacy. This decision can expose the patient to avoidable side effects, compromise therapeutic outcomes, and generate unnecessary expenses.

The transition from childhood to adolescence in these patients represents a critical point for diagnostic reassessment, adaptation of therapeutic strategies, and redefinition of functional goals.

Adolescents with NDDs face increased risks of social isolation, school difficulties, internalizing symptoms, emotional dysregulation, and mental health problems [2,3]. Furthermore, Brazilian health services are rarely prepared to provide continuous and integrated care that encompasses this stage of development [4].

Neurobiological and psychosocial changes in 43 adolescence

Adolescence is marked by significant neurohormonal changes in the brain that occur during puberty, especially in the prefrontal, limbic, and pituitary regions. These changes affect hormonal and emotional regulation, inhibitory control, and decision-making, as well as bodily, genital, and sexual growth [5]. In individuals with NDDs, these changes can exacerbate existing symptoms or contribute to the emergence of new clinical manifestations, such as anxiety, depression, or risky behaviors [6].

From a psychosocial perspective, the challenges of developing autonomy and independence, identity, and sexuality take on distinct contours. Many adolescents with ASD, for example, report difficulties interpreting social norms and establishing emotional bonds [7].

Specificities of neurodevelopmental disorders

Autism spectrum disorder (ASD) 60

During adolescence, individuals with ASD may experience 61 worsening symptoms of irritability, increased stereotypies, 62 or the onset of depression [8]. There is also an increased risk 63

of bullying and social exclusion, which contribute to low 64 self-esteem and greater psychological vulnerability [9].

65

68

69

73

74

75

76

77

79

80

81

88

89

90

93

94

95

96

98

101

102

104

105

107

108

109

110

111

112

113

114

The lack of specialized medical services for adolescents 66 with ASD hinders continuity of care, especially during school 67 and professional transitions [10].

Attention deficit/hyperactivity disorder (ADHD)

Although the core symptoms of ADHD, especially hyperactiv- 70 ity, tend to decrease over time, in up to 60% of cases they 71persist into adolescence and adulthood [11]. Impulsivity and inattention may manifest in a less disruptive manner, but they still have a significant impact on academic performance, social relationships, and mental health [12].

The risk of comorbidities such as mood disorders, substance abuse (e-cigarettes) or alcohol consumption, and risky behaviors (e.g., dangerous challenges) increases during this period [13], requiring continuous monitoring and therapeutic reassessments.

Intellectual disability and other conditions

Adolescents with intellectual disabilities require specific 82 approaches that involve preparation for adulthood, the 83 development of functional skills for some socioeconomic 84 independence, and extended family support [14]. Sexuality and the desire for autonomy also emerge as central topics 86 that need to be discussed sensitively and ethically during consultations with adolescents and their families by pediatricians [15].

Transition models and integrated care

The transition to adolescence should be planned in a struc- 91 tured manner, focusing on the continuity of medical, educational, and social care. Models based on individualized transition plans (ITPs) have proven effective in preparing adolescents for the adult environment [16].

The collaborative work of pediatricians, psychiatrists, educators, and psychotherapists is essential. International 97 guidelines recommend that the transition process ideally begin between the ages of 12 and 14, focusing on developing 99 self-care and decision-making skills [17]. Periodic diagnostic reassessment of these patients is recommended, considering that clinical manifestations and functional demands may change over time, requiring adjustments to therapeutic 103 strategies [18].

Monitoring adolescents over the years shows that in neurodivergent youth, anxiety symptoms appear before puberty and predict the development of depression in cases with autism spectrum disorder, ADHD, suggesting the need for early intervention [19].

At the same time, it is essential to promote the adolescent's progressive autonomy, encouraging their active participation in therapeutic decisions, which contributes to strengthening their identity and engaging in self-care.

The different interventions range from behavioral strategies to psychosocial, pharmacological, and educational support and should be used in an integrated manner. The choice of the ideal therapeutic strategy depends on the adolescent's specific characteristics, their level of functioning, the presence of comorbidities, and the family and social context in which they are inserted.

120

121

122

123

124

125

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160 161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

The focus of therapeutic approaches should be on promoting the individual's overall development and their autonomy in adulthood. Among the main proposed approaches, the authors highlight behavioral therapy, drug management of comorbidities such as anxiety, psychosocial support through support groups, personalized education with pedagogical adaptations and, more recently, the use of assistive technologies such as applications and virtual reality, in addition to continuous family support [20].

Behavioral therapy is often the starting point for interventions targeting neurodivergent patients, especially those with ASD. Applied Behavior Analysis (ABA) has been widely used in this population, with positive results reported in the scientific literature. This approach involves applying principles of behavior analysis to increase desirable behaviors and reduce problematic or avoidant behaviors through positive reinforcement and other behavioral strategies [21].

Furthermore, developing structured routines is a key element of behavioral intervention. Creating clear routines, with set times for study, social activities, and leisure, can help increase a sense of safety and control over one's environment [22].

Although behavioral and psychosocial interventions are considered first-line treatments for neurodivergent individuals, there are cases in which medication is necessary, especially when associated psychiatric comorbidities such as generalized anxiety, depression, or mood disorders are present. In these cases, the judicious use of medications can be essential to improving the adolescent's quality of life and enabling greater engagement in behavioral and educational therapies. Selective serotonin reuptake inhibitors (SSRIs), such as fluoxetine and sertraline, are frequently used to manage anxiety in adolescents [23].

Another essential aspect of therapeutic monitoring for adolescents with a developmental disorder is psychosocial support. Support groups for these adolescents have proven particularly effective in providing a safe and supervised space for these young people to interact with their peers, share experiences, and develop social skills in a structured manner.

Formal education plays a central role in the life of any adolescent. In the case of neurodivergent individuals, adapting the educational environment and curriculum is essential to ensure effective inclusion and academic achievement. The Individualized Educational Plan (IEP) should go beyond basic literacy or behavioral goals and include long-term objectives, such as preparing for university or vocational entrance exams, inclusion in extracurricular activities, and developing functional learning skills for a first job.

It is important that teachers be trained to understand the particularities of these students, adapting their methodologies, using visual aids, concrete language, tasks with welldefined deadlines, and adapted assessments.

Therapies specifically aimed at individuals with ASD have incorporated technological resources that significantly expand the possibilities for intervention. Assistive technologies in the digital world, such as social simulation apps, modeled videos, and interactive games, help adolescents practice social skills in controlled environments with immediate feedback. Many of these apps are specifically developed to promote skills such as emotion recognition, theory of mind development, and understanding of social

182

183

184

185

186

187

188

189

190

191

193

194

195

196

197

198

199

200

203

205

206

207

208

214

215

224

225

228

229

230

233

Another promising resource is virtual reality (VR), which allows simulating complex social situations—such as a job interview, a classroom presentation, or even a meeting with friends—in a completely safe environment. VR enables the replay of experiences, the personalization of scenarios, and performance analysis, all of which contribute to the generalization of learned skills to the real world [25].

It is essential to emphasize the role of the family in the success of any therapeutic plan. Parents and caregivers are 192 not merely observers of the process; they are active participants and often the main mediators of the strategies proposed in therapy. Therefore, family involvement must be systematic and planned.

The transition from adolescence to adulthood in individuals with neurodevelopmental disorders

The transition from adolescence to adulthood marks an important point in human growth, characterized by significant transformations in the biological, psychological, and social domains. For those with neurodevelopmental disorders, this phase often brings additional challenges that 204 require special attention from healthcare professionals, educators, and family members.

Although these disorders begin in early childhood, they often continue throughout life. However, their expressions, needs, and consequences change significantly over time. During adolescence, aspects related to independence, social integration, academic performance, and identity formation become more pronounced. The transition to adulthood brings new functional demands that, in many situations, are not fully met by health and education services that are predominantly focused on children.

The symptoms of NDDs can transform as they transition 216 into adulthood. In ASD, for example, although some repeti- 217 tive behaviors and communication difficulties persist, it is 218 common to observe a refinement in interactional patterns, while new issues such as social anxiety and isolation emerge [26]. In the case of ADHD, longitudinal studies show that hyperactivity tends to decrease with age, while inattention and impulsivity often persist, affecting academic 223 performance and interpersonal relationships [27].

The development of professional and personal autonomy is one of the main goals for people with NDD, and this process is significantly compromised in many cases. Difficulties with emotional self-regulation, time management, and daily tasks directly impact the ability to live independently [28].

Even individuals with good academic and cognitive performance may experience significant deficits in practical 231 daily life skills, such as financial management, personal 232 care, and public transportation.

A study of young adults with ASD showed that only a minority lived independently, held a job, or maintained regular social relationships without additional support [29]. The lack of structured transition programs contributes to this gap, often leaving these young people without support 238 during the most critical phase of their development into adulthood.

240

241

242

243

245 246

247

248

249

250

251

252

253

254 255

256

257 258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

275

276

277

278

279

280 281

282

283

284

285

286

287

288

289

290

291

292

293

294

295 296

297

298

299

300

Permanence in the educational system and the transition to higher education or the job market are particularly challenging stages. Many adolescents with NDD finish high school with curricular adaptations but encounter barriers in continuing their academic careers or finding jobs compatible with their skills [30].

Inclusive education, while a step forward, still shows shortcomings in preparing students with NDD for the demands of adulthood. The lack of individualized plans that address transition goals, such as career guidance, social skills, and independent living practices, is one of the main criticisms raised by experts [31].

In the employment field, young people with NDD face high rates of unemployment or underemployment. In the case of ASD, for example, studies indicate that <20% of autistic adults are able to maintain employment in positions compatible with their cognitive and academic capacities [32]. Barriers related to communication, adaptation to unstructured environments, and difficulties in understanding implicit social rules are factors that hinder these individuals' professional integration.

Regarding specialized support, the transition to adulthood often coincides with the reduction or even cessation of specialized services offered during childhood. Many adolescents with NDD no longer receive regular monitoring by multidisciplinary teams when they reach adulthood (at age 18 and, in many cases, at age 21, according to the Child and Adolescent Statute (ECA), which leads to discontinuity in care and worsening of their clinical condition [33].

Furthermore, mental health issues become more prominent at this stage. Symptoms of depression, suicidal ideation, and self-harm are more common in young adults with NDD than in the general population [34]. Factors such as bullying, social isolation, low self-esteem, and difficulties adapting contribute to this situation of con-

For this reason, it is essential that healthcare systems adopt transition models that ensure continuity of care, with teams prepared to support patients throughout their life cycle. Case management models and coordination between pediatric, adolescent medicine, and adult health services are highlighted as best practices [35].

In this context, families play a central role in the transition to adulthood for young people with NDD. Parents and caregivers often become the primary mediators between the adolescent and the healthcare, education, and employment systems. However, they also face emotional, financial, and physical burdens, especially when there is no effective support network [36].

Cultural and socioeconomic factors also directly influence the autonomy of these young adults. In countries with fewer resources, such as Brazil, or where support services are scarce, the responsibility for care tends to fall entirely on the family, further hindering the development of functional independence [37].

Consequently, public policies should include family support programs, including mental health education, support groups, and legal and financial guidance. Preparation for the future should begin in adolescence, with active parental participation in transition planning.

Several countries have developed specific public policies 301 for the transition of adolescents with chronic conditions and NDD into adulthood. Models such as the UK's "Ready Steady Go" model propose a structured approach, beginning preparation at age 14, with progressive stages leading up to the full transition at age 18 or 21 [38]. This model includes periodic interviews with the young person and their caregivers, the development of individualized plans, and integration with clinical and adult education services.

304

305

306

307

308

309

310

320

321

324

326

328

329

331

332

333

334

335

026

342

343

344

346

347

348

349

351

352

353

356

357

In Brazil, although there are guidelines for care for people with disabilities, such as the 1999 Disability Inclusion 311 Law (Decree N. 3298 of December 20, 1999, which regulates 312 Law N. 7853/89), and establishes the National Policy for the 313 Integration of Persons with Disabilities, consolidating pro- 314 tection standards and establishing guidelines for the social 315 inclusion and citizenship of this population, there are still 316 few services specifically structured for the transition. The lack of coordination between the health, education, and social assistance sectors compromises the continuity of care. The implementation of specialized care teams for adolescents, focusing on life planning, career guidance, and psychosocial support, is urgently needed.

Another relevant aspect of the transition to adulthood is the development of sexuality and romantic relationships. Many adolescents with NDD have difficulty understanding social norms related to intimacy, consent, dating, and even marriage. At the same time, they often have limited access 327 to sex education, due to family fears or the lack of training of health and education professionals [39].

The lack of adequate guidance increases the risk of violence and abuse, sexual exploitation, and inappropriate behavior. Therefore, it is essential that the topic of sexuality be addressed in an accessible, respectful, and ongoing manner. Sex education programs adapted to the language and cognitive abilities of adolescents have proven effective in promoting safety and autonomy [40].

The transition to adulthood for individuals with NDD should not be viewed as a one-off event, but as an ongoing process that requires advance planning, multidisciplinary involvement, and family support. Interventions should begin 340 in adolescence, focusing on the development of practical, 341 social, and emotional skills, as well as strategies for educational and professional inclusion.

Advances in research and public policy formulation have contributed to greater awareness of the needs of this population, but there is still a long way to go. The implementation of structured transition programs, professional training, continuity of mental health care, and support for families is are essential tool to ensure that these adolescent adults reach their potential and lead independent, full, and meaningful lives.

Conclusion

Adolescence for individuals with neurodevelopmental disorders is a phase of multiple challenges and opportunities. Integrated, person-centered, continuous, and collaborative healthcare is essential to promote the well-being and inclusion of these adolescents.

The lack of clear public policies and organized care flows remains a significant obstacle in Brazil. Investments in public 359

4

Jornal de Pediatria xxxx;xxx(xxx): 101463

policies, professional development and training, and integrated transition models are priorities to improve long-term outcomes.

Promoting integration between pediatric services and adolescent mental health professionals is essential to ensure the full development and quality of life of these young people, a human rights issue guaranteed in Article 227 of the Brazilian Federal Constitution.

368 Data availability

- The data that support the findings of this study are available from the corresponding author.
- 371 Conflicts of interest
- 372 The authors declare no conflicts of interest.
- 373 Editor

378

379

380

381

385

386

387

388

389

392

393

394

363

364

365

366

367

374 M.L. Nunes

375 References

- 1. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 5th ed. WA, DC: APA; 2013.
 - Emerson E, Einfeld S. Emotional and behavioural difficulties in young children with and without developmental delay: a binational perspective. J Child Psychol Psychiatry. 2010;51 (5):583–93.
- 3. Doshi-Velez F, Ge Y, Kohane I. Comorbidity clusters in autism spectrum disorders: an electronic health record time-series analysis. Pediatrics. 2014;133(1):e54–63.
 - 4. Viner RM, Ozer EM, Denny S, Marmot M, Resnick M, Fatusi A, et al. Adolescence and the social determinants of health. Lancet. 2012;379(9826):1641–52.
 - 5. Blakemore SJ. Imaging brain development: the adolescent brain. Neuroimage. 2012;61(2):397–406.
- 6. Teliti A, Resulaj L. Adolescence in autism spectrum disorder and challenges encountered. Interdiscip J Res Dev. 2022;9(1):42.
 - 7. Orsmond GI, Kuo HY. The daily lives of adolescents with an autism spectrum disorder: discretionary time use and activity partners. Autism. 2011;15(5):579–99.
- 8. Mazefsky CA, White SW. Emotion regulation: concepts & practice in autism spectrum disorder. Child Adolesc Psychiatr Clin N Am. 2014;23(1):15–24.
- Sterzing PR, Shattuck PT, Narendorf SC, Wagner M, Cooper BP.
 Bullying involvement and autism spectrum disorders: prevalence and correlates of bullying involvement among adolescents with an autism spectrum disorder. Arch Pediatr Adolesc Med.
 2012;166(11):1058-64.
- 403 10. Cheak-Zamora NC, Teti M, First J. Transitions are scary for our
 404 kids, and they're scary for us': family member and youth perspectives on the challenges of transitioning to adulthood with
 406 autism. J Appl Res Intellect Disabil. 2015;28(6):548–60.
- 407 11. Faraone SV, Biederman J. Neurobiology of attention-deficit hyperactivity disorder. Biol Psychiatry. 1998;44(10):951–8.
- 409 12. Cortese S. The neurobiology and genetics of Attention-Deficit/
 410 Hyperactivity Disorder (ADHD): what every clinician should
 411 know. Eur J Paediatr Neurol. 2012;16(5):422-33.

13. Molina BS, Pelham Jr WE. Attention-deficit/hyperactivity disorder and risk of substance use disorder: developmental considerations, potential pathways, and opportunities for research. Annu Rev Clin Psychol. 2014;10:607—39.

414

415

416

417

418

419

420

421

422

423

424

425

426

42.7

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

473

474

475

476

477

478

- 14. Schalock RL, Borthwick-Duffy SA, Bradley VJ, Buntinx WH, Coulter DL, Craig EM, et al. Intellectual disability: definition, classification, and systems of supports. 11th ed. WA, DC: AAIDD; 2010.
- 15. Pownall JD, Jahoda A, Hastings RP. Sexuality and sex education of adolescents with intellectual disability: mothers' attitudes, experiences, and support needs. Intellect Dev Disabil. 2012;50 (2):140–54.
- 16. Rosen DS, Blum RW, Britto M, Sawyer SM, Siegel DM. Society for adolescent medicine. Transition to adult health care for adolescents and young adults with chronic conditions: position paper of the society for adolescent medicine. J Adolesc Health. 2003;33(4):309-11.
- 17. Colver A, Longwell S. New understanding of adolescent brain development: relevance to transitional healthcare for young people with long term conditions. Arch Dis Child. 2013;98(11):902—7.
- 18. Pfeifer JH, Allen NB. Puberty initiates cascading relationships between neurodevelopmental, social, and internalizing processes across adolescence. Biol Psychiatry. 2021;89(2):99–108.
- 19. Orm S, Wood JJ, Fossum IN, Adams K, Andersen PN, Fjermestad K, et al. Anxiety symptoms predict subsequent depressive symptoms in neurodivergent youth: a 10-year longitudinal study. Res Child Adolesc Psychopathol. 2025;53(3):429–41.
- 20. Sociedade Brasileira de Pediatria (SBP). Manual de orientação grupo de trabalho saúde na era digital (gestão 2022-2024). Benefícios da tecnologia para todas as crianças e adolescentes: mais Aptidão, mais Saúde; 2023, [acesso em 2025 jun. 20]. Disponível em: https://www.sbp.com.br/fileadmin/user_upload/23903c-ManOrient_-_BenefTecnol_todas_crc_e_adl-MaisAptidao MaisSaude.pdf.
- 21. Schreibman L. The science and fiction of autism. Cambridge: Harvard University Press; 2005.
- 22. Murray S., Noland B. Video modeling for young children with autism spectrum disorders: a practical guide for parents and professionals. Londres: Jessica Kingsley Publishers; 2013. Williams K, Brignell A, Randall M, Silove N, Hazell P. Selective serotonin reuptake inhibitors (SSRIs) for autism spectrum disorders (ASD). Cochrane Database Syst Rev. 2013;2013(8):CD004677
- 23. Grynszpan O, Weiss PL, Perez-Diaz F, Gal E. Innovative technology-based interventions for autism spectrum disorders: a meta-analysis. Autism. 2014;18(4):346–61.
- 24. Parsons S, Cobb S. State-of-the-art of virtual reality technologies for children on the autism spectrum. Eur J Spec Needs Educ. 2011;26(3):355–66.
- 25. Mandy W, Lai MC. Annual research review: the role of the environment in the developmental psychopathology of autism spectrum condition. J Child Psychol Psychiatry. 2016;57(3):271–92.
- Barkley RA, Murphy KR. Attention-deficit hyperactivity disorder: a clinical workbook. 3rd ed. NY: Guilford Press; 2006.
- 27. Taylor JL, Seltzer MM. Developing a vocational index for adults with autism spectrum disorders. J Autism Dev Disord. 2012;42 (12):2669–79.
- 28. Shattuck PT, Narendorf SC, Cooper B, Sterzing PR, Wagner M, Taylor JL. Postsecondary education and employment among youth with an autism spectrum disorder. Pediatrics. 2012;129 (6):1042–9.
- 29. Newman L, Wagner M, Cameto R, Knokey AM. The post-high school outcomes of youth with disabilities up to 4 years after high school. a report from the national longitudinal transition study-2 (NLTS2). Menlo Park, CA: SRI International; 2011.
- Wehman P, Schall C, McDonough J, Kregel J, Brooke V, Molinelli A, et al. Competitive employment for youth with autism spectrum disorders: early results from a randomized clinical trial. J Autism Dev Disord. 2014;44(3):487–500.

E.J. Silva and E. Eisenstein

479	31. Roux A.M., Rast J.E., Anderson K.A., Shattuck P.T. Nationa
480	autism indicators report: transition into young adulthood. Phila
481	delphia: Life Course Outcomes Research Program; 2015.

- 32. Cheak-Zamora NC, Teti M. "You think it's hard now ... it gets 482 483 much harder for our children": youth with autism and their 484 caregiver's perspectives of health care transition services. Autism. 2015;19(8):992-1001. 485
- 486 33. Hedley D, Uljarević M, Foley KR, Richdale A, Trollor J. Risk and protective factors underlying depression and suicidal ideation 487 488 in autism spectrum disorder. Depress Anxiety. 2018;35(7): 648 - 57. 489
- 34. American Academy of Pediatrics, American Academy of Family 490 Physicians, American College of Physicians, Transitions Clinical 491 492 Report Authoring Group, Cooley WC, Sagerman PJ. Supporting 493 the health care transition from adolescence to adulthood in the medical home. Pediatrics. 2011;128(1):182-200. 494

35. Heller T, Schindler A. Family support interventions for families 495 of adults with intellectual and developmental disabilities. Int 496 Rev Res Ment Retard. 2009;37:299-332.

497

498

499

500

501

502

503

504

506

507

508

509

510

- 36. Schalock RL, Luckasson RA, Shogren KA, Borthwick-Duffy S, Bradley V, Buntinx WH, et al. The renaming of mental retardation: understanding the change to the term intellectual disability. Intellect Dev Disabil. 2007;45(2):116-24.
- 37. NHS England. Ready steady go: transition programme. NHS Foundation Trust; 2018.
- 38. Travers J, Tincani M, Whitby PS, Boutot EA. Alignment of sexuality education with self determination for people with significant 505 disabilities: a review of research and future directions. Educ Train Autism Dev Disabil. 2014;49(2):232-47.
- 39. Ballan MS. Parental perspectives of communication about sexuality in families of children with autism spectrum disorders. J Autism Dev Disord. 2012;42(5):676-84.