



Jornal de
Pediatria

www.jpmed.com.br



ORIGINAL ARTICLE

Serological biomarker models composed of luteinizing hormone, kisspeptin, vitamin D and estradiol, and their clinical test value in girls

Q1 Mingyu Shao^a, Zhiyi Jia^b, Tongtong Liu^b, Xiaopeng Zhang^{b,*}

^a Zibo Central Hospital, Department of Child Health Care, Shandong, China

^b Zibo Central Hospital, Department of Pediatrics, Shandong, China

Received 4 June 2025; accepted 17 October 2025

Available online xxx

KEYWORDS

Q2 Central precocious puberty;
Luteinizing hormone;
Kisspeptin;
Vitamin D;
Estradiol

Abstract

Objective: Central precocious puberty (CPP) may lead to premature pubertal onset. While the GnRH stimulation test remains the gold standard, its invasive nature and prolonged procedure limit clinical utility, particularly in pediatric populations. Emerging biomarkers, including luteinizing hormone, estradiol, kisspeptin, and vitamin D, have shown promise in distinguishing CPP from normal puberty. This study systematically evaluated the diagnostic utility of these biomarkers both individually and in novel combinatorial models, with the aim of establishing a non-invasive and more accessible diagnostic alternative for CPP.

Methods: This retrospective study included 129 girls with CPP and 116 age-matched controls. Clinical characteristics, including height, weight, body mass index, and bone age, were recorded. Serum levels of luteinizing hormone, estradiol, kisspeptin, vitamin D, progesterone, and prolactin were measured. The diagnostic performance of individual biomarkers and three combined biomarker models was assessed using receiver operating characteristic curve analysis.

Results: Model 1 included luteinizing hormone and kisspeptin, Model 2 incorporated vitamin D, and Model 3 added estradiol. CPP patients exhibited significantly higher levels of luteinizing hormone (2.51 vs. 0.23 mIU/mL), kisspeptin (1.59 vs. 0.96 μ g/L), and estradiol (25.86 vs. 13.41 pg/mL) and lower vitamin D levels (20.13 vs. 25.90 ng/mL) compared to controls (all $p < 0.001$). Model 3 demonstrated the highest diagnostic accuracy with an AUC of 0.939 (95% CI: 0.910–0.968), sensitivity of 89.06%, and specificity of 87.93%, outperforming individual biomarkers and other models.

Conclusion: This study highlights the potential of combining luteinizing hormone, kisspeptin, vitamin D, and estradiol into a single diagnostic model for CPP.

© 2025 Published by Elsevier España, S.L.U. on behalf of Sociedade Brasileira de Pediatria. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

* Corresponding author.

E-mail: zhangxiaopeng0322@163.com (X. Zhang).

<https://doi.org/10.1016/j.jpmed.2025.101477>

0021-7557/© 2025 Published by Elsevier España, S.L.U. on behalf of Sociedade Brasileira de Pediatria. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Please cite this article in press as: M. Shao, Z. Jia, T. Liu et al., Serological biomarker models composed of luteinizing hormone, kisspeptin, vitamin D and estradiol, and their clinical test value in girls, *Jornal de Pediatria* (2025), <https://doi.org/10.1016/j.jpmed.2025.101477>

1 Introduction

2 Central precocious puberty (CPP) is a condition marked
3 by the premature activation of the hypothalamic-pitui-
4 tary-gonadal axis, leading to the early onset of secondary
5 sexual characteristics before the age of 8 years in girls
6 and 9 years in boys [1]. CPP is significantly more common
7 in girls and has far-reaching implications for physical,
8 psychosocial, and emotional development, including
9 advanced bone maturation, reduced adult height, and
10 emotional challenges [1,2]. Diagnosing CPP accurately is
11 critical for initiating timely interventions, such as gonad-
12 otropin-releasing hormone (GnRH) analog therapy, which
13 can mitigate long-term consequences of early puberty,
14 and to date, the GnRH stimulation test remains the gold
15 standard method for diagnosing CPP [3]. However, distin-
16 guishing CPP from other variants of early puberty, such
17 as isolated premature thelarche or non-progressive
18 puberty, remains a diagnostic challenge [4].

19 Traditionally, CPP diagnosis has relied on clinical fea-
20 tures, radiographic evidence of advanced bone age, and hor-
21 monal evaluations, including basal and stimulated
22 gonadotropin levels. However, the GnRH stimulation test,
23 although sensitive, is resource-intensive, time-consuming,
24 and inconvenient for pediatric patients [5]. Additionally,
25 gonadotropin levels often overlap between CPP and other
26 conditions, complicating the diagnosis [6]. These limitations
27 highlight the unmet need for non-invasive and reliable bio-
28 markers that can improve diagnostic accuracy. In this con-
29 text, devising a new diagnostic model with higher specificity
30 would be preferred to reduce unnecessary GnRH testing in
31 these benign cases.

32 Recent studies have explored the utility of serum bio-
33 markers such as luteinizing hormone, estradiol, and proges-
34 terone, which are central to the activation of the
35 hypothalamic-pituitary-gonadal axis [7,8]. Emerging bio-
36 markers such as kisspeptin, a neuropeptide regulating gonad-
37 otropin-releasing hormone secretion, have shown promise in
38 distinguishing CPP cases [9]. Additionally, vitamin D, which is
39 inversely associated with puberty progression, offers poten-
40 tial as a diagnostic biomarker [10,11]. However, the diagnos-
41 tic utility of these biomarkers remains limited by variability
42 in sample collection times, pubertal stages, and individual
43 differences [12]. Furthermore, few studies have evaluated
44 combinations of biomarkers to enhance diagnostic accuracy.

45 By leveraging receiver operating characteristic (ROC) curve
46 analysis, researchers can systematically evaluate the diagnos-
47 tic performance of individual biomarkers and biomarker mod-
48 els in terms of sensitivity, specificity, and area under the curve
49 (AUC) [13]. Although individual biomarkers provide important
50 insights, their diagnostic utility is often limited when used in
51 isolation. Integrating biomarkers into combined models has
52 shown potential for improving diagnostic performance, but
53 this approach remains underexplored for CPP diagnosis.

54 This study aims to address these limitations by evaluating
55 the clinical and serological profiles of girls with CPP and
56 assessing the diagnostic value of key biomarkers, including
57 luteinizing hormone, estradiol, vitamin D, and kisspeptin. By
58 constructing and validating three serological biomarker
59 models, this study seeks to identify a non-invasive, efficient,
60 and accurate diagnostic tool for distinguishing CPP from nor-
61 mal puberty. The findings are expected to contribute to the

growing body of evidence supporting biomarker-based diag- 62
nostics for managing CPP [14]. 63

Methods

Study design and participants

This study, being retrospective in nature, was conducted 66
using data from 129 girls diagnosed with central precocious 67
puberty (CPP) who were treated at Zibo Central Hospital. 68
The control group included 116 age-matched girls who 69
underwent routine health examinations during the same 70
period. The diagnostic criteria for ICPP were based on the 71
guidelines referenced from the Chinese Medical Association: 72
(1) Onset of secondary sexual characteristics in girls before 73
age 8 years or menarche occurring under age 10 years; (2) 74
Accelerated linear growth, bone age (BA) exceeding chrono- 75
logical age by 1 year or more; (3) Enlargement of the gonads: 76
pelvic ultrasound shows increased uterine and ovarian vol- 77
ume in girls, with multiple ovarian follicles > 4 mm in diam- 78
eter; (4) Activation of the HPGA: The serum levels of 79
gonadotropins and sex hormones reached pubertal levels. 80
Exclusion criteria comprised: (1) Secondary central preco- 81
cious puberty with a clearly defined organic cause; (2) Partic- 82
ipants taking known medications affecting the HPGA or 83
having used steroid medications prior to the study; (3) 84
Peripheral precocious puberty; (4) Known endocrine diseases 85
or chromosomal abnormalities were excluded from the 86
study; (5) Severe organic diseases and incomplete medical 87
records; (6) Individuals with corneal inflammation, blephari- 88
tis, conjunctival stones, entropion, and abnormal blinking 89
due to local eye diseases and ocular irritation factors were 90
excluded; (7) Those with abnormal blinking resulting from 91
refractive errors, drugs, psychological, and systemic ill- 92
nesses were also excluded; (8) Individuals with a history of 93
eye trauma, surgery, or corneal contact lens wear were not 94
included. The participant selection process is outlined in the 95
study flowchart. The study was approved by Zibo Central 96
Hospital, the approval number was 2010-SMY. The study was 97
performed in strict accordance with the Declaration of Hel- 98
sinki, Ethical Principles for Medical Research Involving 99
Human Subjects. Written informed consent was waived due 100
to the nature of the retrospective analysis. 101

Clinical and biomarker measurements

Baseline clinical characteristics, including height, weight, 103
body mass index (BMI), and bone age, were recorded for all 104
participants. Blood samples were collected, and serum lev- 105
els of basal luteinizing hormone, progesterone, estradiol, 106
prolactin, vitamin D, and kisspeptin were measured using 107
standardized assays. The concentrations of these biomarkers 108
were compared between the CPP and control groups. 109

Biomarker model development and evaluation

ROC curve analysis was conducted to evaluate the diagnostic 111
performance of individual biomarkers, including luteinizing 112
hormone, estradiol, vitamin D, and kisspeptin, in distin- 113
guishing CPP cases from controls. Serological biomarker 114
models were developed to assess combined diagnostic 115

utility. Model 1 included luteinizing hormone and kisspeptin, Model 2 incorporated luteinizing hormone, kisspeptin, and vitamin D, and Model 3 included luteinizing hormone, kisspeptin, vitamin D, and estradiol. ROC curves were generated for each model, and diagnostic performance was assessed by calculating the area under the curve (AUC), 95 % confidence intervals (CI), sensitivity, specificity, and cut-off values. These parameters were compared to determine the diagnostic utility of each model. To verify the scientific validity of the study design, the selection of this specific combination was based on step-wise logistic regression.

Statistical analysis

Continuous variables were expressed as medians with interquartile ranges, and categorical variables as frequencies and percentages. Between-group comparisons were performed using the Mann-Whitney U test for continuous data and the chi-squared test for categorical data. ROC curve analysis was used to evaluate the diagnostic performance of individual biomarkers and combined models, and AUC values were compared using statistical software. Statistical significance was set at $p < 0.05$. All analyses were performed using SPSS.

Results

Baseline characteristics indicate accelerated growth and maturation in CPP

A total of 245 girls were included in the study, with 116 in the control group and 129 in the CPP group. The participant selection and group allocation are detailed in Figure 1. The baseline clinical characteristics, shown in Table 1, revealed no significant difference in chronological age between the two groups ($p = 0.991$). However, girls in the CPP group exhibited significantly higher height ($p < 0.001$), weight ($p < 0.001$), BMI ($p = 0.039$), and bone age ($p < 0.001$), indicating accelerated growth and skeletal maturation compared to the control group. These findings indicate that CPP is associated with distinct physical development patterns that can help differentiate it from normal puberty.

Elevated luteinizing hormone, kisspeptin, and estradiol characterize CPP

To further explore the hormonal differences underlying these physical changes, serum biomarker analysis was performed. The CPP group exhibited significantly higher levels of




Figure 1 Flow chart of the study participants.

Table 1 Clinical characteristics at baseline of the girls in the two groups.

Items	Control group (n = 116)	CPP group (n = 129)	p-Value
chronological age (years)	7.0 (7.0–8.0)	7.0 (7.0–8.0)	0.991
Height (cm)	123.00 (120.00–125.50)	130.70 (124.70–136.15)	<0.001
Weight (kg)	24.20 (20.90–26.10)	27.30 (23.90–31.80)	<0.001
BMI (kg/m ²)	16.10 (14.47–17.05)	16.30 (15.44–17.65)	0.039
bone age (years)	7.64 (7.10–8.20)	9.64 (8.65–10.13)	<0.001

Values of $p < 0.05$ were considered statistically significant. BMI: body mass index.

Table 2 Serum levels of luteinizing hormone, progesterone, prolactin, kisspeptin, vitamin D and estradiol in the two groups.

Items	Control group (n = 116)	CPP group (n = 129)	p-value
Luteinizing hormone (mIU/mL)	0.23 (0.08–0.45)	2.51 (0.69–3.84)	<0.001
Progesterone (ng/mL)	0.36 (0.21–0.73)	0.46 (0.26–0.87)	0.135
Prolactin (ng/mL)	9.75 (7.11–11.98)	8.89 (5.55–18.33)	0.814
Kisspeptin ($\mu\text{g/L}$)	0.96 (0.52–1.49)	1.59 (0.53–2.39)	<0.001
Vitamin D (ng/mL)	25.90 (22.64–31.17)	20.13 (16.94–24.11)	<0.001
Estradiol (pg/mL)	13.41 (9.86–16.98)	25.86 (18.45–33.18)	<0.001

Values of $p < 0.05$ were considered statistically significant.

luteinizing hormone (2.51 mIU/mL vs. 0.23 mIU/mL, $p < 0.001$), kisspeptin (1.59 $\mu\text{g/L}$ vs. 0.96 $\mu\text{g/L}$, $p < 0.001$), and estradiol (25.86 pg/mL vs. 13.41 pg/mL, $p < 0.001$) compared to the control group (Table 2). In contrast, vitamin D levels were significantly lower in the CPP group (20.13 ng/mL vs. 25.90 ng/mL, $p < 0.001$), suggesting an inverse relationship between vitamin D and markers of CPP. No significant differences were observed in progesterone and prolactin levels between the two groups. These biomarker patterns reflect distinct endocrine profiles in girls with CPP, with elevated gonadotropins and sex hormones as defining characteristics.

Diagnostic performance of biomarkers is enhanced by combining multiple indicators

Building on these biomarker differences, ROC curve analysis was conducted to evaluate the diagnostic utility of individual biomarkers and their combined models. Among the individual biomarkers, luteinizing hormone and estradiol demonstrated the highest diagnostic accuracy (Figure. 2A). Recognizing the potential for improved performance, three biomarker models were developed by combining various indicators. Model 1, which included luteinizing hormone and




Figure 2 ROC curves for the control group and the CPP group and the combined auxiliary diagnostic models. (A) ROC curves for luteinizing hormone, kisspeptin, vitamin D and estradiol. (B) ROC curve for model 1. (C) ROC curve for model 2. (D) ROC curve for model 3. ROC, receiver operating characteristic.

kisspeptin, achieved an AUC of 0.784 (95 % CI: 0.726–0.842) with a sensitivity of 73.44 % and specificity of 75.86 %. Adding vitamin D in Model 2 further improved diagnostic performance, achieving an AUC of 0.902 (95 % CI: 0.864–0.940), sensitivity of 81.25 %, and specificity of 86.21 % (Figure. 2B–2C). Model 3, which incorporated estradiol in addition to the other biomarkers, exhibited the highest diagnostic performance with an AUC of 0.939 (95 % CI: 0.910–0.968), sensitivity of 89.06 %, and specificity of 87.93 % (Figure. 2D). These results demonstrate that combining multiple biomarkers significantly enhances diagnostic precision for CPP.

Combining biomarkers provides the best diagnostic utility for CPP

A comparison of the diagnostic models is presented in Table 3. While Model 1 provided moderate diagnostic accuracy, the inclusion of vitamin D in Model 2 resulted in a notable improvement in both sensitivity and specificity. Model 3, which added estradiol to the biomarker panel, outperformed the other models, offering the most accurate tool for CPP diagnosis. In addition, since the two indicators, progesterone and prolactin, showed no significant difference, they were not included in the final model. These findings highlight the value of integrating multiple biomarkers into diagnostic models, emphasizing their utility in clinical settings for improving the diagnosis of CPP. The results demonstrate that Model 3 represents a non-invasive, efficient, and highly accurate diagnostic approach, offering a promising alternative to traditional methods.

Discussion

This study evaluated the diagnostic utility of key serum biomarkers and their combinations in distinguishing CPP from normal puberty. The present findings reveal significant differences in clinical and biochemical profiles between the CPP and control groups, highlighting the potential of integrating biomarkers into diagnostic models. These results address the limitations of traditional methods, such as the GnRH stimulation test, which, while effective, is invasive, time-consuming, and often impractical in routine clinical settings [4,5].

The clinical characteristics observed in this study, including higher height, weight, BMI, and bone age in the CPP group, align with previous research documenting

accelerated growth and skeletal maturation in CPP patients [1,8]. However, while these characteristics support clinical suspicion of CPP, their specificity is limited due to overlap with other conditions such as premature thelarche or non-progressive puberty [6]. This underscores the importance of incorporating reliable biomarkers into diagnostic frameworks to improve accuracy and specificity.

Among the biomarkers analyzed, luteinizing hormone, kisspeptin, and estradiol were significantly elevated in girls with CPP, while vitamin D levels were notably reduced. These findings are consistent with earlier studies highlighting the role of these hormones in pubertal activation and progression [7,9]. Kisspeptin, a central regulator of pubertal onset, demonstrated particular diagnostic potential, consistent with previous studies showing its elevated levels in CPP cases [10]. Vitamin D, inversely associated with pubertal progression, provides additional diagnostic insight to this study, further validating earlier research suggesting its association with delayed puberty onset [11,15]. The inclusion of vitamin D in biomarker models represents an underexplored area that enhances the diagnostic framework presented in this study.

The ROC curve analysis revealed the limitations of individual biomarkers when used in isolation. Luteinizing hormone and estradiol demonstrated moderate diagnostic accuracy, consistent with earlier studies reporting AUC values of 0.7–0.8 for these markers [8,16]. However, combining biomarkers into models significantly improved diagnostic performance. Model 3, which included luteinizing hormone, kisspeptin, vitamin D, and estradiol, achieved the highest accuracy with an AUC of 0.939, surpassing most previously reported single-marker analyses. The diagnostic improvements achieved by integrating biomarkers align with emerging evidence suggesting that biomarker combinations provide greater sensitivity and specificity compared to single markers [13]. Although the current study has indicated a cut-off value for Model 3 of 0.770, caution should be taken when applying this threshold in guiding similar studies, given limitations such as the retrospective nature and single-center design of the current study. Currently, this model may serve as a screening tool for GnRH testing - individuals with high scores should undergo GnRH testing for CPP diagnosis, while those with low scores may be suitable for clinical observation. Future large-scale studies will be conducted to evaluate whether this model could replace the gold standard GnRH test.

Table 3 Characteristics of the auxiliary diagnostic models.

Model	Included indicators	AUC	95 % CI	p-value	Sensitivity (%)	Specificity (%)	Cut-off value
Model 1	luteinizing hormone, kisspeptin	0.784	0.726–0.842	<0.001	73.44	75.86	0.493
Model 2	luteinizing hormone, kisspeptin, vitamin D	0.902	0.864–0.940	<0.001	81.25	86.21	0.675
Model 3	luteinizing hormone, kisspeptin, vitamin D, estradiol	0.939	0.910–0.968	<0.001	89.06	87.93	0.770

Values of $p < 0.05$ were considered statistically significant.

AUC, area under the curve; CI, confidence interval.

Since the two indicators, progesterone and prolactin, showed no significant difference, they were not included in the final model.

Compared to previous research, this study provides a more comprehensive evaluation of integrated biomarker models. While earlier studies focused on traditional markers like luteinizing hormone and estradiol, this study incorporated kisspeptin and vitamin D, demonstrating their combined utility in enhancing diagnostic accuracy [10]. By addressing gaps in prior research, this study offers a novel and practical approach for diagnosing CPP.

Despite the promising results, certain limitations should be acknowledged. The retrospective nature of this study may introduce biases related to data collection and selection criteria [17]. In addition, the single-center design may limit the generalizability of these findings. Larger, multicenter studies are needed to validate these biomarker models across diverse populations. Furthermore, the cost-effectiveness and clinical feasibility of implementing these models in routine practice remain areas for future investigation [18].

Conclusions

In conclusion, this study highlights the value of combining multiple biomarkers, including luteinizing hormone, kisspeptin, vitamin D, and estradiol, to improve the accuracy of CPP diagnosis. Model 3, which demonstrated superior sensitivity and specificity, offers a robust and non-invasive diagnostic tool for CPP. By incorporating a broader range of biomarkers and integrating them into clinically relevant models, this study addresses limitations in previous research and advances the field of biomarker-based diagnostics for CPP [19]. Future research should focus on validating these models in larger cohorts and assessing their implementation in clinical practice to optimize the diagnosis and management of CPP [20,21].

Funding

None.

Statement of ethics

The study was approved by Zibo Central Hospital; the approval number was 2010-SMY. The study was performed in strict accordance with the Declaration of Helsinki, Ethical Principles for Medical Research Involving Human Subjects.

Informed consent

The need for informed consent was waived by Zibo Central Hospital.

Data availability

The data that support the findings of this study are not publicly available due to privacy reasons, but are available from Xiaopeng Zhang upon reasonable request.

Conflicts of interest

The authors declare no conflicts of interest.

Editor

C. de A.D Alves

References

- [1]. Carel JC, Eugster EA, Rogol A, Ghizzoni L, Palmert MR, et al. Consensus statement on the use of gonadotropin-releasing hormone analogs in children. ESPE-LWPES GnRH analogs consensus conference group. *Pediatrics*. 2009;123(4):e752–62.
- [2]. Cohen A, Dempster DW, Recker RR, Lappe JM, Zhou H, Zwanlen A, et al. Abdominal fat is associated with lower bone formation and inferior bone quality in healthy premenopausal women: a transiliac bone biopsy study. *J Clin Endocrinol Metab*. 2013;98(6):2562–72.
- [3]. Pellegrin MC, Marzin C, Monasta L, Tamaro G, Vidonis V, Vittori G, et al. A short-duration gonadotropin-releasing hormone stimulation test for the diagnosis of central precocious puberty. *Medicina*. 2023;60(1):24.
- [4]. Vargas KM, Da Cunha C, Andreolini R. Amphetamine and pentylenetetrazole given post-trial 1 enhance one-trial tolerance to the anxiolytic effect of diazepam in the elevated plus-maze in mice. *Prog Neuropsychopharmacol Biol Psychiatry*. 2006;30(8):1394–402.
- [5]. Ludvigsson J, Krisky D, Casas R, Battelino T, Castano L, Greening J, et al. GAD65 antigen therapy in recently diagnosed type 1 diabetes mellitus. *N Engl J Med*. 2012;366(5):433–42.
- [6]. Mulligan KJ, McCauley H. Factors that influence the determination of residual solvents in pharmaceuticals by automated static headspace sampling coupled to capillary GC–MS. *J Chromatogr Sci*. 1995;33(1):49–54.
- [7]. Greulich WW, Pyle SI. Radiographic atlas of skeletal development of the hand and wrist. 2nd Edition Stanford: Stanford University Press; 1959. <https://doi.org/10.1097/00000441-195909000-00030>.
- [8]. Parent AS, Teilmann G, Juul A, Skakkebaek NE, Toppari J, Bourguignon JP. The timing of normal puberty and the age limits of sexual precocity: variations around the world, secular trends, and changes after migration. *Endocr Rev*. 2003;24(5):668–93.
- [9]. Pinilla L, Aguilar E, Dieguez C, Millar RP, Tena-Sempere M. Kisspeptins and reproduction: physiological roles and regulatory mechanisms. *Physiol Rev*. 2012;92(3):1235–316.
- [10]. Lee JM, Smith JR, Philipp BL, Chen TC, Mathieu J, Holick MF. Vitamin D deficiency in a healthy group of mothers and newborn infants. *Clin Pediatr*. 2007;46(1):42–4.
- [11]. Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007;357(3):266–81.
- [12]. Szendroedi J, Phielix E, Roden M. The role of mitochondria in insulin resistance and type 2 diabetes mellitus. *Nat Rev Endocrinol*. 2011;8(2):92–103.
- [13]. Obuchowski NA. Receiver operating characteristic curves and their use in radiology. *Radiology*. 2003;229(1):3–8.
- [14]. Truzzi JC, Gomes CM, Bezerra CA, Plata IM, Campos J, Garrido GL, et al. Overactive bladder - 18 years - part II. *Int Braz J Urol*. 2016;42(2):199–214.
- [15]. Misra M, Pacaud D, Petryk A, Collett-Solberg PF, Kappy M. Drug and therapeutics committee of the lawson wilkins pediatric endocrine society. vitamin D deficiency in children and its management: review of current knowledge and recommendations. *Pediatrics*. 2008;122(2):398–417.

- 370 [16]. Kapellen TM, Reimann R, Kiess W, Kostev K. Prevalence of 383
371 medically treated children with ADHD and type 1 diabetes in 384
372 Germany - analysis of two representative databases. *J Pediatr* 385
373 *Endocrinol Metab.* 2016;29(11):1293–7. 386
- 374 [17]. Ghaedi E, Mohammadi M, Mohammadi H, Ramezani-Jolfaie N, 387
375 Malekzadeh J, Hosseinzadeh M, et al. Effects of a paleolithic 388
376 diet on cardiovascular disease risk factors: a systematic 389
377 review and meta-analysis of randomized controlled trials. *Adv* 390
378 *Nutr.* 2019;10(4):634–46. Erratum in: *Adv Nutr.* 2020;11 391
379 (4):1054. 392
- 380 [18]. Pinna G, Hiedra L, Meinhold H, Eravci M, Prengel H, Brödel O, 393
381 et al. 3,3'-Diiodothyronine concentrations in the sera of 394
382 patients with nonthyroidal illnesses and brain tumors and of 395
healthy subjects during acute stress. *J Clin Endocrinol Metab.* 383
1998;83(9):3071–7. 384
- [19]. Butler TR, Self RL, Smith KJ, Sharrett-Field LJ, Berry JN, Littleton 385
JM, et al. Selective vulnerability of hippocampal cornu ammonis 386
1 pyramidal cells to excitotoxic insult is associated with the 387
expression of polyamine-sensitive N-methyl-D-aspartate-type glu- 388
tamate receptors. *Neuroscience.* 2010;165(2):525–34. 389
- [20]. Malátová I, Fojtík P. Activity of ¹³⁷Cs in human body of inhabi- 390
tants of the Czech republic. *Radiat Prot Dosimetry.* 2019;186 391
(2–3):196–201. 392
- [21]. Nematollahi A, Rezai H, Helan JA, Moghaddam N. Occurrence 393
of *linguistula serrata* nymphs in cattle slaughtered in Tabriz. 394
Iran J Parasit Dis. 2015;39(2):140–3. 395